

Density of states “width parity” effect in d -wave superconducting quantum wires

K. Ziegler¹, W.A. Atkinson^{2,3}, P.J. Hirschfeld²

¹Institut für Physik, Universität Augsburg, 86135 Augsburg, Germany

²Department of Physics, University of Florida, PO Box 118440, Gainesville FL 32611 USA

³Department of Physics, Southern Illinois University, Carbondale, IL 62901 USA.

(February 1, 2008)

We calculate the density of states (DOS) in a clean mesoscopic d -wave superconducting quantum wire, i.e. a sample of infinite length but finite width N . For open boundary conditions and half-filling, the DOS at zero energy is found to be zero if N is even, and nonzero if N is odd. At finite chemical potential, all chains are gapped but the qualitative differences between even and odd N remain.

I. INTRODUCTION

In recent years the fabrication of nanoscale devices has heightened interest in solid-state quantum systems where the discreteness of the level spacing plays an important role. Originally discussed only in semiconductor quantum dots and quantum wires, following the work of Ralph et al [1] many fascinating consequences of the level discreteness were also observed in ultrasmall metal particles. These included the observation of a spectroscopic gap attributed to pair correlations in the small grains which depended on the so-called “number parity” of the sample, i.e. whether the number of electrons in the grain was even or odd. The physics of the levels in such grains has been quite well understood. [2] Superconducting systems extended in one dimension but mesoscopic in another have also been studied. [3]

In most of these studies, a gap with A_{1g} or “s-wave” symmetry has been assumed for simplicity, unless otherwise stated. However, for some practical purposes it may be of interest to use cuprate samples, which are known to have $d_{x^2-y^2}$ symmetry. From a fundamental point of view it is also expected that these systems will be different because already in the bulk state there are states below the maximum gap scale; the pure d -wave bulk DOS is linear in energy, $\rho(E) \sim |E|$. We have studied the DOS of mesoscopic d -wave quantum wires and found a new kind of “parity effect”, not related to the number of particles in the system, but to the parity of the number of “chains” across the mesoscopic sample. In some ways, the effect is reminiscent of a type of mesoscopic gap effect which has been recently discussed in the context of single-wall carbon nanotubes, where the existence or nonexistence of a gap depends on the intrinsic twist or chirality induced while forming the tube. [5] The d -wave systems we discuss, while they may be considered to be “tubes” if periodic boundary conditions are employed, do not break either time reversal symmetry or spatial parity. Nevertheless, at half-filling, we observe that if the width of the system N is even there is a gap in the DOS, whereas if the width is odd the density of states at zero energy $\rho(0)$ is nonzero. As N is increased, both the even- N gap and the

odd- N $\rho(0)$ vanish as they must to give the bulk d -wave result $\rho(E) \sim |E|$. Although the DOS vanishes at $E = 0$ for all N when $\mu \neq 0$, there is still a pronounced even-odd effect in the DOS, and we argue that this effect should be observable in quantum wires fabricated from cuprate superconductors. We speculate further that it may be of relevance to the study of disorder-induced pseudogaps in bulk d -wave superconductors, which have been the subject of much controversy in recent years. [4]

II. MODEL

We consider the DOS of a d -wave superconducting chain (DWSC) that is coupled to $N - 1$ other DWSCs. The chains form a 2D system, and the first chain is at the boundary. Similar problems were considered in previous studies of random flux systems [8] and quasi-one-dimensional disordered tight-binding models [9,10]. In this paper we will ignore the effect of disorder and study the odd-even effect in a pure system of coupled DWSCs. Such a behavior is also known for a system of coupled spin chains which has a gap (no gap) if the number of chains is even (odd) [6,7].

There are several possible geometries one can assume for this problem. The most natural, and the one we consider here, is a wire parallel to the 100 (or 010) crystal direction. In this geometry, the boundaries of the wire are aligned with both the underlying CuO_2 crystal and the gap maxima (which are tied to the crystal lattice). Other geometries, such as a 110 oriented wire, are expected to show similar even-odd effects to the ones described below, but these will be mixed with Andreev scattering resonances [11] and thus will be more difficult to interpret. In addition, we anticipate that the 100 oriented wire is the most accessible to current technology.

The Bogoliubov-de Gennes Hamiltonian of the 2D DWS

$$H = (-\nabla^2 + \mu)\sigma_3 + \Delta\sigma_1$$

is defined on a square lattice with

$$-\nabla^2 f(r) = -t \sum_{j=1}^2 [f(r + e_j) + f(r - e_j)]$$

and the DWS order parameter

$$\Delta f(r) = \Delta_0 \sum_{j=1}^2 (-1)^j [f(r + e_j) + f(r - e_j)].$$

With this definition, the gap in familiar k -space representation is $\Delta_k = 2\Delta_0[\cos(k_x) - \cos(k_y)]$. In the following calculations we will measure energies in units of t for simplicity. Moreover, it is assumed that chains are infinitely long and arranged parallel to the y -axis of our 2D system.

III. PERIODIC BOUNDARIES

The finite set of chains can be closed periodically in the x -direction by identifying the first with the $N + 1^{\text{th}}$ chain. While this is not particularly physical, the calculation is fairly transparent and therefore worthwhile for pedagogical reasons. For this translational-invariant N -chain system the integration in y -direction (i.e. along the chains) can be performed and gives the DOS at zero energy

$$\begin{aligned} \rho(E=0) &= -\frac{1}{\pi N} \sum_{k_x} \int_{-\pi}^{\pi} \frac{dk_y}{2\pi} \text{Im } G_0(\mathbf{k}, +i\epsilon) \\ &= \lim_{\epsilon \rightarrow 0} \frac{\epsilon \rho_0}{N} \sum_{n=0}^{N-1} \frac{1}{\sqrt{\epsilon^2 + [\mu + 4t \cos(k_x)]^2}}, \end{aligned} \quad (1)$$

where

$$\rho_0 = \frac{2t}{\pi \sqrt{t^2 + \Delta_0^2} \sqrt{16t^2 - \mu^2}}.$$

Here G_0 is the Green's function for a clean 2D d -wave superconductor, $k_x = 2\pi n/N$ ($n = 0, 1, \dots, N-1$), and we have set the lattice constant to unity. First, we consider the DOS at half-filling ($\mu = 0$). We find that $\rho(E=0)$ is *finite* for N a multiple of 4 since the allowed values of k_x include $k_x = \pi/2$ and $3\pi/2$ which produce finite contributions to Eq. (1). For other values of N , on the other hand, expression (1) always vanishes. At $\mu = 0$, then, we have

$$\rho_N(E=0) = \begin{cases} 2\rho_0/N & \text{for } N = 4k, k = 1, 2, \dots \\ 0 & \text{otherwise} \end{cases}.$$

This periodicity in N reflects the conditions necessary to form a standing wave in the x -direction at the Fermi energy. For $\mu = 0$, it is the four nodal points at $\mathbf{k} = (\pm\pi/2, \pm\pi/2)$ which produce the standing wave. The wave function at $E = 0$ is thus exactly the same as in the

normal state; it is therefore not surprising that $\rho(E=0)$ is only weakly dependent on Δ_0 .

For given N , there will be a discrete set of μ which produce a finite density of states at $E = 0$. For generic μ , the nodal eigenstates do not contribute at $E = 0$, but we expect on the basis of (1) that a qualitative width parity effect will remain at nonzero E . As we shall see in the next section, numerical studies on realistic models find a pronounced effect of this type.

IV. OPEN BOUNDARIES

Periodic boundary conditions are not realistic for typical experiments because the superconductor is a planar object. Therefore, a better choice is open boundaries if we have a situation with a finite number of chains. The DOS is calculated numerically for $N = 4, 5, 20, 21$, as shown in Figs. 1-5. Since we have just shown that Δ_0 plays a minor role in the parity effect, we proceed (for simplicity) under the assumption $\Delta_0/t = 1$ unless explicitly stated otherwise. In numerical work (e.g. Fig. 6 below), it is shown that this assumption does not change the qualitative results. From the figures the alternating structure of the DOS for odd and even number of chains is immediately obvious, and may be characterized by the DOS of the end chain.

A. Recursive method

Some insight into the dependence of these gaps and residual DOS's on system size and other parameters may be obtained from analytical methods. First, the Hamiltonian H can be diagonalized with respect to y -direction by a Fourier transformation $y \rightarrow k_y$. Then the structure with respect to the x -direction can be expressed in a matrix form for $x = 1, 2, \dots, N$ as

$$\hat{H}_N = \begin{pmatrix} H_{NN} & H_{NN-1} & 0 & \dots & 0 \\ H_{N-1N} & H_{N-1N-1} & \ddots & \ddots & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & H_{22} & H_{21} \\ 0 & \dots & 0 & H_{12} & H_{11} \end{pmatrix}.$$

For a compact notation we introduce a block matrix representation

$$\hat{H}_N := \begin{pmatrix} H_{NN} & \hat{H}_{NN-1} \\ \hat{H}_{N-1N} & \hat{H}_{N-1} \end{pmatrix}$$

with

$$\hat{H}_{NN-1} = (H_{NN-1}, 0, \dots, 0), \quad (2)$$

$$\hat{H}_{N-1N} = \begin{pmatrix} H_{N-1N} \\ 0 \\ \vdots \\ 0 \end{pmatrix}. \quad (3)$$




FIG. 1. Density of states for four and five chains with $\Delta_0/t = 1$ and $\mu = 0$.

The corresponding Green's function at zero energy, which is required for the evaluation of the DOS, can then be written (after a shift $H \rightarrow H + i\epsilon\sigma_0$) as

$$\begin{aligned} [\hat{H}_N^{-1}]_{NN} &= (H_{NN} - \hat{H}_{NN-1}\hat{H}_{N-1}^{-1}\hat{H}_{N-1N})^{-1} \\ &= (H_{NN} - H_{NN-1}[\hat{H}_{N-1}^{-1}]_{N-1N-1}H_{N-1N})^{-1}. \end{aligned}$$

The second equation is due to the special form of the matrices given in (2) and (3). Using $\Gamma_N = [\hat{H}_N^{-1}]_{NN}$ the recurrence relation reads

$$\Gamma_N = (H_{NN} - H_{NN-1}\Gamma_{N-1}H_{N-1N})^{-1}. \quad (4)$$

This is a “continued fraction representation” of the 2×2 matrix Γ_N , the Green's function of end chain labeled by N at a given k_y . It can be used even in the presence of random terms along the x -direction because no diagonalization of the matrix is necessary.

Now we will apply the recursive method to evaluate the DOS of the end chain of N DWSCs. The calculation is simplified greatly if we assume $\Delta_0 = t = 1$ (the more




FIG. 2. Density of states for 4 chains: DOS of the end chain and average over all chains, with $\Delta_0/t = 1$ and $\mu = 0$.

general case is treated in numerical calculations). In this special case we have

$$\begin{aligned} H_{NN} &= \begin{pmatrix} \mu + 2 \cos k_y + i\epsilon & -2 \cos k_y \\ -2 \cos k_y & -\mu - 2 \cos k_y + i\epsilon \end{pmatrix} \\ &= (\mu + 2 \cos k_y)\sigma_3 - 2 \cos k_y\sigma_1 + i\epsilon\sigma_0 \\ &\equiv h + i\epsilon\sigma_0 \end{aligned} \quad (5)$$

and

$$H_{NN-1} = H_{N-1N} = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} = \sigma_1 + \sigma_3.$$

The initial value (single chain) is

$$\Gamma_1 = (H_{11})^{-1} = \frac{-i\epsilon\sigma_0 + (\mu + 2 \cos k_y)\sigma_3 - 2 \cos k_y\sigma_1}{\epsilon^2 + (\mu + 2 \cos k_y)^2 + 4 \cos^2 k_y}.$$

B. Few Chains

By direct iteration of Eq. (4) we can obtain the DOS analytically for the end chain of a system of N chains,





FIG. 3. Density of states for 5 chains: DOS of the end chain and average over all chains, with $\Delta_0/t = 1$ and $\mu = 0$.

provided we restrict consideration to $\mu = 0$. Using $c = 2 \cos k_y$, for $N = 1$ we get

$$\rho_1(0) = \lim_{\epsilon \rightarrow 0} \frac{\epsilon}{\pi} \int_{-\pi}^{\pi} \frac{1}{\epsilon^2 + 2c^2} \frac{dk_y}{2\pi} = \frac{1}{\sqrt{8}\pi}$$

in agreement with the result in Sect. III. For $N = 2$ we have

$$\rho_2(0) = \lim_{\epsilon \rightarrow 0} \frac{\epsilon}{\pi} \int_{-\pi}^{\pi} \frac{1}{\epsilon^2 + 2c^2 + 2} \frac{dk_y}{2\pi}.$$

From the pole structure of the integrand, we see that there is a gap $2E_g = 2\sqrt{2} \approx 2.83$. In the case of $N = 3$ there is again a nonzero DOS at $E = 0$

$$\rho_3(0) = \lim_{\epsilon \rightarrow 0} \frac{\epsilon}{\pi} \int_{-\pi}^{\pi} \frac{\epsilon^2 + 2c^2 + 2}{(\epsilon^2 + 2c^2)^2 + 4(\epsilon^2 + 2c^2)} \frac{dk_y}{2\pi} = \frac{1}{2\sqrt{8}\pi}.$$

and for $N = 4$

$$\rho_4(0) = \lim_{\epsilon \rightarrow 0} \frac{\epsilon}{\pi} \int_{-\pi}^{\pi} \frac{\epsilon^2 + 2c^2 + 4}{(\epsilon^2 + 2c^2)^2 + 6(\epsilon^2 + 2c^2) + 4} \frac{dk_y}{2\pi},$$

which has a gap $2E_g = 2\sqrt{3 - \sqrt{5}} \approx 1.75$ (see Fig. 1). In general the energy gap observed depends on the size of the superconducting order parameter maximum Δ_0 in a nonlinear way, although relatively simple expressions may be obtained for a small number of chains.

The reader will notice that the even-odd effect described here differs from the periodic boundary case, where gapless states arise for N a multiple of 4. This is expected, since the effect we are reporting arises from self-interference of quantum wavefunctions in a nanoscale system. The path-length for a constructively interfering closed path in the x -direction is $2L = 2n\pi/k_F$ for open boundary conditions, and $L = 2n\pi/k_F$ for periodic boundary conditions.





FIG. 4. Density of states for 20 chains: DOS of the end chain and average over all chains, with $\Delta_0/t = 1$ and $\mu = 0$.

C. Many Chains: Stationary Behavior for $N \rightarrow \infty$

While in systems mesoscopic in both directions (finite length L quantum wires), even-odd parity effects in N are known to survive the thermodynamic limit $L \rightarrow \infty$




FIG. 5. Density of states for 21 chains: DOS of the end chain and average over all chains, with $\Delta_0/t = 1$ and $\mu = 0$.

[10], they must disappear as $N \rightarrow \infty$ when we recover the fully 2D system. This is evident from the numerical evaluation of the DOS for large N (Figures 4 and 5), and we can see this analytically by showing that as $N \rightarrow \infty$ $\rho(0) = 0$ independent of N .

A single iteration of Eq. (4) yields a relation between the Green's functions of an even (or odd) number of chains, respectively

$$\Gamma_N = (H_{NN} - H_{NN-1}[H_{N-1N-1} - H_{N-1N-2}\Gamma_{N-2}H_{N-2N-1}]^{-1}H_{N-1N})^{-1}. \quad (6)$$

For the DWSCs this reads

$$\Gamma_N = (H_{NN} - [\frac{1}{2}(\sigma_1 + \sigma_3)H_{N-1N-1}(\sigma_1 + \sigma_3) - \Gamma_{N-2}]^{-1})^{-1}.$$

For constant $\mu = 0$ we get from expression (5)

$$H_{N-1N-1} = h + i\epsilon\sigma_0$$

and

$$\frac{1}{2}(\sigma_1 + \sigma_3)H_{N-1N-1}(\sigma_1 + \sigma_3) = -h + i\epsilon\sigma_0$$

such that

$$\Gamma_N = (h + i\epsilon\sigma_0 + [h - i\epsilon\sigma_0 + \Gamma_{N-2}]^{-1})^{-1}$$

with the initial expressions

$$\Gamma_1 = \frac{1}{\epsilon^2 + 2c^2}(h - i\epsilon\sigma_0)$$

and

$$\Gamma_2 = (1 + \frac{2}{\epsilon^2 + 2c^2})^{-1}\Gamma_1.$$

The symmetric 2×2 matrix h can be diagonalized by a orthogonal transformation, leading to diagonal elements $\lambda_{1/2} = \pm\sqrt{2}c$. This also diagonalizes the initial expressions $\Gamma_{1,2}$ as well as the recurrence relation

$$\gamma_{N+2,j} = \frac{1}{\lambda_j + i\epsilon + \frac{1}{\lambda_j - i\epsilon + \gamma_{N,j}}} \quad (7)$$

with diagonal elements $\gamma_{N,1/2}$ of Γ_N . Then the recursion (7) has two fixed points (i.e., stationary states)

$$\bar{\gamma}_j^\pm = -\frac{z_j^*}{2}(1 \pm \sqrt{1 + 4/|z_j|^2})$$

with $z_j = \lambda_j + i\epsilon = -(-1)^j\sqrt{2}c + i\epsilon$. For $\epsilon > 0$ a positive imaginary part of $\gamma_{N,j}^+$ implies a positive imaginary part of $\gamma_{N+2,j}^+$. Since the imaginary parts of the initial values are positive, only the fixed point $\bar{\gamma}_j^+$ can be reached in the case under consideration.

The DOS of the end chain can be calculated from the fixed point and reads

$$\rho(0) = -\frac{1}{\pi} \lim_{\epsilon \rightarrow 0} \text{Im} \int_{-\pi}^{\pi} (\bar{\gamma}_1^+ + \bar{\gamma}_2^+) \frac{dk_y}{2\pi} \propto \lim_{\epsilon \rightarrow 0} \epsilon \ln(1/\epsilon) = 0.$$

This reflects the well-known result of the infinite 2D DWSC which has a linear pseudogap.

D. Realistic models

The preceding discussion has assumed a half-filled tight-binding band and $\Delta_0/t = 1$; both conditions correspond to special symmetries of the Hamiltonian which might be expected to influence the form of the spectrum. The assumption $\Delta_0/t = 1$ was in fact made simply in order to obtain the analytical results discussed in sections IV A-C, and it may easily be checked numerically (Figure 6) that for small values of Δ_0/t and $\mu = 0$, the qualitative even-odd effect in the chain width parity is still obtained. This is consistent with the qualitative results in the case of periodic boundaries.

More significant changes occur when $\mu \neq 0$. In Figure 7 we see that for an even number of chains, the full gap in all chains is preserved, while for an odd number, those chains which at half-filling had a finite density of states




FIG. 6. Density of states for four and five chains with $\Delta_0/t = 0.3$ and $\mu = 0$.

have now acquired a small gap. Analytic results for this case are complicated and not particularly enlightening, but it is clear that for $\mu \ll \Delta_0$, the small gap on the x =odd chains in the odd N case is of the order of μ itself. This is also evident by analogy to Eq. (1), where the pole in the denominator of the integral is shifted by the chemical potential.

A more realistic parameter choice for the optimally doped cuprates would correspond to a doping of 15%, or about $\mu \simeq 0.3t$ for the simple tight-binding spectrum on a square lattice, and a much smaller gap magnitude, of order $\Delta_0 = 0.1t$. In this case the large “gap” is set by $2\Delta_0$, but in the odd-chain systems there is a much smaller gap on alternating chains, in Figure 8. Thus the density of states still generally exhibits a pronounced odd-even width parity effect.

V. CONCLUSIONS

We have exhibited, using both analytical and numerical means, a number parity effect in the width N




FIG. 7. Density of states for four and five chains with $\Delta_0/t = 1$ and $\mu/t = 0.3$.

of a mesoscopic d -wave superconducting quantum wire, wherein a finite DOS is found at the Fermi level for odd N , and zero DOS (with energy gap) for even N . The result should be of some practical interest in the not-so-distant future for nanoscale d -wave wire structures. Superconductivity in single layers of the cuprates has already been demonstrated, and it seems plausible that a sample of controlled width might be fabricated, and that an STM experiment on such a sample would be able to observe the effects we predict.

It is interesting to note some unusual features of the system we treat here, to our knowledge for the first time, in the presence of impurities. First of all, although the superconducting order parameter has d -wave symmetry, the system may (for N even) have a full gap in the excitation spectrum. In such a situation, one may ask what is the effect of isolated impurities added to the system, and at least naively would obtain single-impurity d -wave like bound states in this gap without broadening arising from coupling to the quasiparticle continuum. Since the density of states in the odd- N systems has an oscillatory behavior across the sample width, the existence and life-




FIG. 8. Density of states for four and five chains with $\Delta_0/t = 0.1$ and $\mu/t = 0.3$.

time of these states is expected to depend sensitively on their location in the wire as well.

In addition, the observation we make here may be of interest to the study of the influence of disorder on the DOS of fully 2D d -wave superconductors, a subject which has received intense attention recently. [4] The result appears to be that the DOS is generically zero in a 2D d -wave superconductor at zero energy, but can be constant or divergent in cases manifesting special symmetries. At the same time, Brouwer et al. [10] have shown in disordered quasi1D systems with chiral symmetry that $\rho(0)$ is zero or divergent according to the number parity of the wire width. It appears that the states in odd or even chain systems are modified in different ways by the combined effects of level repulsion and symmetry. Understanding the differences in the odd or even approach to the thermodynamic limit in the presence of disorder may give insight into the physics of localization and DOS suppression in the 2D d -wave case, which is still poorly understood. We will address these questions in a subsequent work.

Acknowledgements

This work is supported by NSF grants DMR-9974396

and INT-9815833, and an exchange grant from DAAD. The authors are grateful to J. von Delft and C. Mudry for useful discussions. PH is grateful to the Institute for Theoretical Physics for its hospitality during work on the manuscript.

-
- [1] D.C. Ralph, C.T. Black, and M. Tinkham, PRL 74 3241 (1995).
 - [2] F. Braun and J. von Delft, Phys. Rev. Lett. 81, 4712 (1998).
 - [3] A.D. Zaikin, D.S. Gobulev, A. van Otterlo, and G.T. Zimanyi, Phys. Rev. Lett. 78, 1552 (1997).
 - [4] A. A. Nersisyan, A. M. Tsvelik, and F. Wenger, Nucl. Phys. B **438**, 561 (1995); Phys. Rev. Lett. **72**, 2628 (1994); K. Ziegler, M.H. Hettler, and P.J. Hirschfeld, Phys. Rev. Lett. 77, 3013 (1996); Phys. Rev. B 57, 10825 (1998); T. Senthil, Matthew P. A. Fisher, Leon Balents, and Chetan Nayak, Phys. Rev. Lett. **81** 4704 (1998); T. Senthil and Matthew P. A. Fisher, Phys. Rev. B60, 6893 (1999); C. Pépin and P.A. Lee, Phys. Rev. B **63** 54502 (2001); W.A. Atkinson, P.J. Hirschfeld, A. H. MacDonald, and K. Ziegler, Phys. Rev. Lett. **85**, 3926 (2000).
 - [5] A. Kleiner and S. Eggert, cond-mat/0007244.
 - [6] F.D.M. Haldane, Phys. Rev. Lett. 61, 1029 (1988).
 - [7] E. Fradkin, Field Theory of Condensed Matter Systems, Addison-Wesley, Redwood City (1991).
 - [8] J. Miller and J. Wang, Phys. Rev. Lett. **76**, 1461 (1996).
 - [9] P.W. Brouwer, C. Mudry, B.D. Simons, and A. Altland, Phys. Rev. Lett. **81**, 862 (1998).
 - [10] P.W. Brouwer, C. Mudry, and A. Furusaki, Phys. Rev. Lett. 84, 2913 (2000).
 - [11] M. Fogelstrom, D. Rainer, and J. A. Sauls, Phys. Rev. Lett. **79**, 281 (1997).